

HIGHLIGHTS OF PRESCRIBING INFORMATION
These highlights do not include all the information needed to use **ROPIVACINE HYDROCHLORIDE INJECTION, safely and effectively. See full prescribing information for ROPIVACINE HYDROCHLORIDE INJECTION.**
• **Signs of methemoglobinemia may occur.** (5.4)
ROPIVACINE HYDROCHLORIDE INJECTION, for epidural, perineural, or infiltration use
U.S. APPROX: 1996

INDICATIONS AND USE
Roipivacine Hydrochloride injection is an amide local anesthetic indicated in adults for the production of local or regional anesthesia for surgery and for acute pain management. **Surgical Anesthesia:** epidural block for surgery including cesarean section; major nerve block; local infiltration. **Acute Pain Management:** epidural continuous infusion or intermittent bolus, e.g., postoperative or labor; local infiltration.

DOSE AND ADMINISTRATION
• See Table 1 for Dosage Recommendations (2.2).

DOSE FORMS AND STRENGTHS
• Injection: 2 mg/mL (0.2%) 5 mg/mL (0.5%), or 10 mg/mL (1%) in single-dose vials (3)
• Injection: 2 mg/mL (0.2%) in single-dose, ready-to-use, polypropylene flexible bag overwrapped with aluminum pouch (3).

CONTRAINDICATIONS
History of hypersensitivity to local anesthetics of the amide type (4).

FULL PRESCRIBING INFORMATION: CONTENTS*
1. **INDICATIONS AND USAGE**
2. **DOSE AND ADMINISTRATION**
3. **DOSE FORMS AND STRENGTHS**
4. **CONTRAINDICATIONS**
5. **WARNINGS AND PRECAUTIONS**
5.1 General Warnings and Precautions
5.2 Unintended Intravenous Injection
5.3 Intra-Arterial Infusions and Risk of Chondrolysis
5.4 Risk of Methemoglobinemia
5.5 Central Nervous System Toxicity
5.6 Epidural Anesthesia
5.7 Use in Brachial Plexus Block
5.8 Use in Peripheral Nerve Block
5.9 Use in Head and Neck Area
5.10 Use in Ophthalmic Surgery
5.11 Hepatic Disease
5.12 Cardiovascular Impairment
5.13 Risk of Additive Effects
5.14 Malignant Hyperthermia
6. **ADVERSE REACTIONS**
7. **DRUG INTERACTIONS**

WARNINGS AND PRECAUTIONS
• Delay in proper management of dose-related toxicity, underventilation, and/or altered sensitivity may lead to the development of acidosis, cardiac arrest and, possibly, death. (5.1)
• In performing Ropivacine hydrochloride blocks, unintended intravenous injection is possible and may result in cardiac arrhythmia or cardiac arrest. (5.2)
• Intra-arterial infusions of local anesthetics may cause chondrolysis. Ropivacine hydrochloride is not approved for this use. (5.3)
• Signs of methemoglobinemia may occur. (5.4)

ADVERSE REACTIONS
• Most common adverse reactions (incidence $\geq 1\%$) are hypotension, nausea, vomiting, bradycardia, fever, pain, postoperative complications, anemia, paresthesia, headache, pruritus, and back pain. (6)
To report SUSPECTED ADVERSE REACTIONS, contact Caplin Steriles Limited at 1-866-978-9111 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch

DRUG INTERACTIONS
• Agents structurally related to amide-type local anesthetics: Concurrent use may cause additive effects. (7)
See 17 for PATIENT COUNSELING INFORMATION.

USE IN SPECIFIC POPULATIONS
8.1 Pregnancy
8.2 Lactation
8.3 Pediatric Use
8.4 Geriatric Use
8.5 Hepatic Impairment
8.6 Renal Impairment

DESCRIPTION
11.1 Description
11.2 Mechanism of Action
11.3 Pharmacodynamics
11.4 Pharmacokinetics
11.5 Clinical Pharmacology

NONCLINICAL TOXICOLOGY
13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility
13.2 Animal Toxicology and/or Pharmacology
13.3 Human Toxicology

CLINICAL STUDIES
14.1 Epidural Administration in Surgery
14.2 Epidural Administration in Cesarean Section
14.3 Epidural Administration in Labor and Delivery
14.4 Epidural Administration in Postoperative Pain Management
14.5 Peripheral Nerve Block
14.6 Local Infiltration

HOW SUPPLIED/STORAGE AND HANDLING
16.1 How Supplied
16.2 Storage and Handling

PATIENT COUNSELING INFORMATION
17.1 Information for Patients and Caregivers

***Sections or subsections omitted from the full prescribing information are not listed.**

Revised: 09/2024

FULL PRESCRIBING INFORMATION

1. INDICATIONS AND USAGE

Ropivacine Hydrochloride is indicated for the production of local or regional anesthesia for surgery and for acute pain management.

Surgical Anesthesia: epidural block for surgery including cesarean section; major nerve block; local infiltration.

Acute Pain Management: epidural continuous infusion or intermittent bolus, e.g., postoperative or labor; local infiltration.

2. DOSE AND ADMINISTRATION

There have been adverse event reports of chondrolysis in patients receiving intra-articular infusions of local anesthetics following arthroscopic and other surgical procedures. Ropivacine hydrochloride is not approved for this use [see **Warnings and Precautions (5.3)**].

The rapid injection of a large volume of local anesthetic solution should be avoided and fractional (incremental) doses should always be used. The smallest dose and concentration required to produce the desired result should be administered.

The dose of any local anesthetic administered varies with the anesthetic procedure, the area to be anesthetized, the vascularity of the tissues, the number of neuronal segments to be blocked, the depth of anesthesia and degree of muscle relaxation required, the duration of anesthesia desired, individual tolerance, and the physical condition of the patient. Patients in poor general condition due to aging or other compromising factors such as partial or complete heart conduction block, advanced liver disease or severe renal dysfunction require special attention although regional anesthesia is frequently indicated in these patients. To reduce the risk of potentially serious adverse reactions, attempts should be made to optimize the patient's condition before major blocks are performed, and the dosage should be adjusted accordingly.

Use an adequate test dose (2 to 5 mL of a short acting local anesthetic solution containing epinephrine) prior to induction of complete block. This test dose should be repeated if the patient is moved in such a fashion as to have displaced the epidural catheter. Allow adequate time for onset of anesthesia following administration of each test dose.

These products are intended for single use and are not for use in preservatives. Any solution remaining from an opened container should be discarded promptly. In addition, continuous infusion bottles should not be left in place for more than 24 hours.

2.2 Dosage Recommendations

Table 1 Dosage Recommendations						
	Conc. mg/mL	(%)	Volume mL	Dose mg	Onset min	Duration hours
SURGICAL ANESTHESIA						
Lumbar Epidural	0.5	(0.5%)	15 to 30	7.5 to 150	15 to 30	2 to 4
Administration	7.5	(0.75%)	15 to 25	113 to 188	10 to 20	2 to 5
Surgery	10	(1%)	15 to 20	150 to 200	10 to 20	4 to 6
Lumbar Epidural	0.5	(0.5%)	20 to 30	100 to 150	15 to 25	2 to 4
Administration	7.5	(0.75%)	15 to 20	113 to 150	10 to 20	2 to 5
Cesarean Section	0.5	(0.5%)	5 to 15	2.5 to 7.5	10 to 30	n/a ^a
Thoracic Epidural	0.5	(0.5%)	5 to 15	38 to 113	10 to 30	n/a ^a
Surgery	7.5	(0.75%)	10 to 15	113 to 226	10 to 20	n/a ^a
Major Nerve Block ^b (e.g., brachial plexus block)	0.5	(0.5%)	35 to 50	175 to 250	15 to 30	5 to 6
Field Block	0.5	(0.5%)	10 to 40	75 to 300	10 to 25	4 to 10
(e.g., minor nerve blocks and infiltration)	0.5	(0.5%)	1 to 20	5 to 200	1 to 15	2 to 6
LABOR PAIN MANAGEMENT						
Lumbar Epidural Administration	0.2	(0.2%)	10 to 20	2 to 20	10 to 15	0.5 to 1.5
Continuous infusion ^c	2	(0.2%)	6 to 14 mL/h	12 to 28 mg/h	n/a ^a	n/a ^a
Incremental infusions (top-up) ^d	2	(0.2%)	10 to 15 mL	20 to 30 mg	n/a ^a	n/a ^a
POSTOPERATIVE PAIN MANAGEMENT						
Lumbar Epidural Administration	0.2	(0.2%)	6 to 14 mL/h	12 to 28 mg/h	n/a ^a	n/a ^a
Continuous infusion ^c	2	(0.2%)	6 to 14 mL/h	12 to 28 mg/h	n/a ^a	n/a ^a
Thoracic Epidural Administration	0.2	(0.2%)	6 to 14 mL/h	12 to 28 mg/h	n/a ^a	n/a ^a
Continuous infusion ^c	2	(0.2%)	1 to 10 mL/h	5 to 200 mg/h	1 to 5	2 to 6
(e.g., minor nerve block)	5	(0.5%)	1 to 40 mL/h	5 to 200 mg/h	1 to 5	2 to 6

^a – Not Applicable

1 – The dose for a major nerve block must be adjusted according to site of administration and patient status. Suprascavicular brachial plexus blocks may be associated with a higher frequency of serious adverse reactions, regardless of the local anesthetic used [see **Warnings and Precautions (5.7)**].

2 – Median dose of 2 mg per hour was administered by continuous infusion or incremental injections (top-up) over a median delivery time of 5.5 hours.

3 – Cumulative dose up to 770 mg of Ropivacine Hydrochloride over 24 hours (intrathecal plus postoperative infusion).

Continuous epidural infusion at rates up to 28 mg per hour for 72 hours were well tolerated in adults, i.e., 2016 mg plus surgical dose of approximately 100 to 150 mg as top-up.

The doses in the table are those considered to be appropriate to produce a successful block and should be regarded as guidelines for use in adults. Individual variations in onset and duration occur. The figures reflect the expected average dose range needed. For other local anesthetic techniques standard current textbooks should be consulted.

5.7 Use in Brachial Plexus Block

Ropivacine plasma concentrations may approach the threshold for central nervous system toxicity after the administration of 300 mg of ropivacaine for brachial plexus block. Caution should be exercised when using the 300 mg dose [see **Overdose (10)**].

The dose for a major nerve block must be adjusted according to the site of administration and patient status. Suprascavicular brachial plexus blocks may be associated with a higher frequency of serious adverse reactions, regardless of the local anesthetic used.

5.8 Use in Peripheral Nerve Block

Major peripheral nerve blocks may result in the administration of a large volume of local anesthetic in highly vascularized areas, often close to large vessels where there is an increased risk of intravascular injection and/or rapid systemic absorption, which can lead to high plasma concentrations.

5.9 Use in Head and Neck Area

Small doses of local anesthetics injected into the head and neck area may produce adverse reactions similar to systemic toxicity seen with unintentional intravascular injections of larger doses. The injection procedures require the utmost care. Confusion, convulsions, respiratory depression, and/or respiratory arrest, and cardiovascular stimulation or depression have been reported. These reactions may be due to intra-arterial injection of the local anesthetic with retrograde flow to the cerebral circulation. Patients receiving these blocks should have their circulation and respiration monitored and be constantly observed. Resuscitative equipment and personnel for treating adverse reactions should be immediately available. Dosage recommendations should not be exceeded [see **Dosage and Administration (2.2)**].

5.10 Use in Ophthalmic Surgery

The use of Ropivacine hydrochloride in retrobulbar blocks for ophthalmic surgery has not been studied. Until appropriate experience is gained, the use of Ropivacine hydrochloride for such surgery is not recommended.

5.11 Hepatic Disease

Because amide-type local anesthetics such as ropivacaine are metabolized by the liver, these drugs, especially repeat doses, should be used cautiously in patients with hepatic disease. Patients with severe hepatic disease, because of their inability to metabolize local anesthetics normally, are at a greater risk of developing toxic plasma concentrations.

5.12 Cardiovascular Impairment

Ropivacine should also be used with caution in patients with impaired cardiovascular function because they may be less able to compensate for functional changes associated with the prolongation of A-V conduction produced by these drugs.

5.13 Risk of Additive Effects

Ropivacine hydrochloride should be used with caution in patients receiving other local anesthetics or agents structurally related to amide-type local anesthetics, since the toxic effects of these drugs are additive. [See **Drug Interactions (7)**].

Patients treated with class II antiarrhythmic drugs (e.g., amiodarone) should be under close surveillance and ECG monitoring considered, since cardiac effects may be additive. [See **Drug Interactions (7)**].

5.14 Malignant Hyperthermia

Many drugs used during the conduct of anesthesia are considered potential triggering agents for malignant hyperthermia (MH). Amide-type local anesthetics are not known to trigger this reaction. However, since the need for supplemental general anesthesia cannot be predicted in advance, it is suggested that a standard protocol for MH management should be available.

6. ADVERSE REACTIONS

Reactions to ropivacaine are characteristic of those associated with other amide-type local anesthetics. A major cause of adverse reactions to this group of drugs may be associated with excessive plasma levels, which may be due to overdose, unintentional intravascular injection or slow metabolic degradation.

The reported adverse events are derived from clinical studies conducted in the U.S. and other countries. The reference drug was usually bupivacaine. The studies used a variety of premedications, sedatives, and surgical procedures of varying length. A total of 3,988 patients have been exposed to Ropivacine hydrochloride at concentrations (as percentage) that occurred in at least 1% of Ropivacine hydrochloride-treated patients in these studies. The majority of patients receiving concentrations higher than 5 mg/mL (0.5%) were treated with Ropivacine hydrochloride. Because clinical trials are conducted under widely conditions, adverse reactions rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Incidence $\geq 5\%$

For the indications of epidural administration in surgery, cesarean section, postoperative pain management, peripheral nerve block, and local infiltration, the following treatment-emergent adverse events were reported with an incidence of $\geq 5\%$ in all clinical studies (N=3988): hypotension (37%), nausea (24.8%), vomiting (11.6%), bradycardia (9.3%), fever (8.2%), pain (8%), postoperative complications (7.1%), anemia (6.1%), paresthesia (5.6%), headache (5.1%), pruritus (5.1%), and back pain (5%).

Incidence 1 to 5%

Urinary retention, dizziness, rigors, hypertension, tachycardia, anxiety, oliguria, hyposthesia, chest pain, hypokalemia, dyspnea, cramps, and urinary tract infection.

Incidence in Controlled Clinical Trials

The reported adverse events are derived from controlled clinical studies with Ropivacine hydrochloride (concentrations ranged from 0.125% to 1.5% for Ropivacine hydrochloride and 0.25% to 0.75% for bupivacaine) in the U.S. and other countries involving 3,094 patients. Table 2 and Table 3 list adverse events reported in patients receiving ropivacaine at concentrations of 0.5 mg/mL (0.5%) or 1.0 mg/mL (1.0%).

Hypotension	536	(32.3)	408	(28.5)
Nausea	283	(17)	207	(14.4)
Vomiting	117	(7)	86	(6.1)
Bradycardia	96	(5.8)	73	(5.1)
Headache	84	(5.1)	68	(4.7)
Paresthesia	82	(4.9)	57	(4)
Back pain	73	(4.4)	75	(5.2)
Pain	71	(4.3)	71	(5)
Pruritus	63	(3.8)	40	(2.8)
Fever	61	(3.7)	37	(2.6)
Dizziness	42	(2.5)	23	(1.6)
Rigors (Chills)	42	(2.5)	24	(1.7)
Post operative complications	41	(2.5)	44	(3.1)
Hypoesthesia	27	(1.6)	24	(1.7)
Urinary retention	23	(1.4)	20	(1.4)
Progression of labor poor/tailed	23	(1.4)	22	(1.5)
Anxiety	21	(1.3)	11	(0.8)
Breast disorder, breast-feeding	21	(1.3)	12	(0.8)
Rhinitis	18	(1.1)	13	(0.9)

Table 3 Adverse Events Reported in ≥1% of Fetuses or Neonates of Mothers Who Received Regional Anesthesia (Cesarean Section and Labor Studies)				
Adverse Reaction	Ropivacaine hydrochloride total N=639		Bupivacaine total N=573	
	N	%	N	%
Fetal bradycardia	77	(12.1)	68	(11.9)
Neonatal jaundice	49	(7.7)	47	(8.2)
Neonatal complication-NOS	42	(6.6)	38	(6.6)
Apgar score low	18	(2.8)	14	(2.4)
Neonatal respiratory disorder	17	(2.7)	18	(3.1)
Neonatal tachypnea	14	(2.2)	15	(2.6)
Neonatal fever	13	(2)	14	(2.4)
Fetal bradycardia	13	(2)	12	(2.1)
Fetal distress	11	(1.7)	10	(1.7)
Neonatal infection	10	(1.6)	8	(1.4)
Neonatal hypoglycemia	8	(1.3)	6	(2.8)

Incidence $<1\%$

The following adverse events were reported during the Ropivacine hydrochloride clinical program in more than one patient (N=3988), occurred at an overall incidence of $<1\%$, and were considered relevant:

Application Site Reactions – injection site pain	Cardiovascular System – vasovagal reaction, syncope, postural hypotension, non-specific ECG abnormalities
Female Reproductive – poor progression of labor, uterine atony	Gastrointestinal System – local incontinence, ileus, nausea, retching, vomiting
General and Other Disorders – hypothermia, malaise, asthenia, accident and/or injury	Hearing and Vestibular – tinnitus, hearing abnormalities
Heart Rate and Rhythm – extrasystoles, non-specific arrhythmia, atrial fibrillation	Metabolic Disorders – hypoglycemia
Life and Dilatory System – jaundice	Musculoskeletal System – myalgia
Neurological System – ST segment changes, myocardial infarction	Nervous System – tremor, Horner's syndrome, paresthesia, dykinesia, neuroprathy, vertigo, coma, convulsion, hypokinesia, hypotonia, phosic, stupor
Psychiatric Disorders – agitation, confusion, somnolence, nervousness, amnesia, hallucination, emotional lability, insomnia, nightmares	Respiratory System – bronchospasm, coughing
Skin Disorders – rash, urticarial	Urinary System Disorders – urinary incontinence, micturition disorder
Vascular – deep vein thrombosis, phlebitis, pulmonary embolism	Vision – vision abnormalities

For the indication epidural anesthesia for surgery, the 15 most common adverse events were compared between different concentrations of Ropivacine hydrochloride and bupivacaine. Table 4 is based on data from trials in the U.S. and other countries where Ropivacine hydrochloride was administered as an epidural anesthetic for surgery.

Adverse Reaction	Ropivacine hydrochloride				Bupivacaine			
	5 mg/mL		7.5 mg/mL		10 mg/mL		5 mg/mL	
	total N=256		total N=297		total N=207		total N=236	
	N	%	N	%	N	%	N	%
hypotension	99	(38.7)	146	(49.2)	113	(54.6)	91	(38.6)
nausea	34	(13.3)	68	(22.9)	47	(22.7)	36	(15.3)
bradycardia	29	(11.3)	58	(19.5)	40	(19.3)	32	(13.6)

back pain	18	(7)	23	(7.7)	34	(14.4)	21	(8.9)	23	(13.2)
vomiting	18	(7)	33	(11.1)	23	(11.1)	19	(8.1)	14	(8)
headache	12	(4.7)	20	(6.7)	16	(7.7)	13	(5.5)	9	(5.2)
fever	8	(3.1)	5	(1.7)	18	(8.7)	11	(4.7)		
chills	6	(2.3)	7	(2.4)	6	(2.9)	4	(1.7)	3	(1.7)
urinary retention	5	(2)	8	(2.7)	10	(4.8)	10	(4.2)		
paresthesia	5	(2)	10	(3.4)	5	(2.4)	7	(3)		
pruritus			14	(4.7)	3	(1.4)			7	(4)

Using data from the same studies, the number (%) of patients experiencing hypotension is displayed by patient age, drug and concentration in Table 5. In Table 6, the adverse events for Ropivacine hydrochloride are broken down by gender.

Table 5 Effects of Age on Hypotension (Epidural Administration)										
Total N: Ropivacaine hydrochloride = 760, Bupivacaine = 410										
	Ropivacaine hydrochloride						Bupivacaine			
AGE	5 mg/mL		7.5 mg/mL		10 mg/mL		5 mg/mL		7.5 mg/mL	
	N	%	N	%	N	%	N	%	N	%
< 65	68	(32.2)	99	(43.2)	87	(51.5)	64	(33.5)	73	(48.3)
≥65	31	(68.9)	47	(69.1)	26	(68.4)	27	(66.0)	16	(69.6)

**Table 6 Most Common Adverse Events by Gender (Epidural Administration)
Total N: Females = 405, Males = 355**

Adverse Reaction	Female		Male	
	N	(%)	N	(%)
hypotension	220	(54.3)	138	(38.9)
nausea	119	(29.4)	23	(6.5)
bradycardia	65	(16)	56	(15.8)
vomiting	59	(14.6)	8	(2.3)
back pain	41	(10.1)	23	(6.5)
headache	33	(8.1)	17	(4.8)
chills	18	(4.4)	5	(1.4)
fever	16	(4)	3	(0.8)
pruritus	16	(4)	1	(0.3)
pain	12	(3)	4	(1.1)
urinary retention	11	(2.7)	7	(2)
dizziness	9	(2.2)	4	(1.1)
hypoaesthesia	8	(2)	2	(0.6)
paresthesia	8	(2)	10	(2.8)

Risk Summary

One publication reported that ropivacaine is present in human milk at low levels following administration of ropivacaine in women undergoing cesarean section. No adverse reactions were reported in the infants. There is no available information on the drug's effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for Ropivacaine hydrochloride and any potential adverse effects on the breastfed child from Ropivacaine hydrochloride or from the underlying maternal condition.

8.4 Pediatric Use

The safety and efficacy of Ropivacaine hydrochloride in pediatric patients have not been established.

8.5 Geriatric Use

Of the 2,978 subjects that were administered Ropivacaine hydrochloride injection in 71 controlled and uncontrolled clinical studies, 803 patients (27%) were 65 years of age or older which includes 127 patients (4%) 75 years of age and over. Ropivacaine hydrochloride injection was found to be safe and effective in the patients in these studies. Clinical data in one published article indicate that differences in various pharmacodynamic measures were observed with increasing age. In one study, the upper level of analgesia increased with age, the maximum decrease of mean arterial pressure (MAP) declined with age during the first hour after epidural administration, and the intensity of motor blockade increased with age.

This drug and its metabolites are known to be excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Elderly patients are more likely to have decreased hepatic, renal, or cardiac function, as well as concomitant disease. Therefore, care should be taken in dose selection, starting at the low end of the dosage range, and it may be useful to monitor renal function [see Clinical Pharmacology (12.3)].

8.6 Hepatic Impairment

Because amide-type local anesthetics such as ropivacaine are metabolized by the liver, these drugs, especially repeat doses, should be used cautiously in patients with hepatic disease. Patients with severe hepatic disease, because of their inability to metabolize local anesthetics normally, are at a greater risk of developing toxic plasma concentrations [see Warnings and Precautions (5.1)].

8.7 Renal Impairment

This drug and its metabolites are known to be excreted by the kidney, and the risk of toxic reactions to this drug may be greater in patients with impaired renal function. Therefore, care should be taken in dose selection, starting at the low end of the dosage range, and it may be useful to monitor renal function [see Clinical Pharmacology (12.3)].

10 OVERDOSAGE

Acute emergencies from local anesthetics are generally related to high plasma levels encountered, or large doses administered, during therapeutic use of local anesthetics or to unintended subarachnoid or intravascular injection of local anesthetic solution [see Adverse Reactions (6) and Warnings and Precautions (5.1, 5.2, 5.6)].

10.1 Treatment

Therapy with Ropivacaine hydrochloride should be discontinued at the first sign of toxicity. No specific information is available for the treatment of toxicity with Ropivacaine hydrochloride; therefore, treatment should be symptomatic and supportive. The first consideration is prevention, best accomplished by incremental injection of Ropivacaine hydrochloride, careful and constant monitoring of cardiovascular and respiratory vital signs and the patient's state of consciousness after each local anesthetic and during continuous infusion. At the first sign of change in mental status, oxygen should be administered.

The first step in the management of systemic toxic reactions, as well as underventilation or apnea due to unintentional subarachnoid injection of drug solution, consists of immediate attention to the establishment and maintenance of a patent airway and effective assisted or controlled ventilation with 100% oxygen with a delivery system capable of permitting immediate positive airway pressure by mask. Circulation should be assisted as necessary. This may prevent convulsions if they have not already occurred.

If necessary, use drugs to control convulsions. Intravenous barbiturates, anticonvulsant agents, or muscle relaxants should only be administered by those familiar with their use. Immediately after the institution of these ventilatory measures, the adequacy of the circulation should be evaluated. Supportive treatment of circulatory depression may require administration of intravenous fluids, and, when appropriate, a vasopressor dictated by the clinical situation (such as ephedrine or epinephrine to enhance myocardial contractile force).

Should cardiac arrest occur, prolonged resuscitative efforts may be required to improve the probability of a successful outcome.

The mean dosages of ropivacaine producing seizures, after intravenous infusion in dogs, nonpregnant and pregnant sheep were 4.9, 6.1 and 5.9 mg/kg, respectively. These doses were associated with peak arterial total plasma concentrations of 11.4, 4.3 and 5 mg/mL, respectively.

In human volunteers given intravenous Ropivacaine hydrochloride, the mean (min-max) maximum tolerated total and free arterial plasma concentrations were 4.3 (3.4 to 5.3) and 0.6 (0.3 to 0.9) mcg/mL, respectively, at which time moderate CNS symptoms (muscle twitching) were noted.

Clinical data from patients experiencing local anesthetic induced convulsions demonstrated rapid development of hypoxia, hypercarbia and acidosis within a minute of the onset of convulsions. These observations suggest that oxygen consumption and carbon dioxide production are greatly increased during local anesthetic convulsions and emphasize the importance of immediate and effective ventilation with oxygen which may avoid cardiac arrest.


If difficulty is encountered in the maintenance of a patent airway or if prolonged ventilatory support (assisted or controlled) is indicated, endotracheal intubation, employing drugs and techniques familiar to the clinician, may be indicated after initial administration of oxygen by mask.

The same position is dangerous in pregnant women at term because of aorticocaval compression by the gravid uterus. Therefore, during treatment of systemic toxicity, maternal hypotension or fetal bradycardia following regional block, the parturient should be maintained in the left lateral decubitus position if possible, or manual displacement of the uterus off the great vessels should be accomplished. Resuscitation of obstetrical patients may take longer than resuscitation of non-pregnant patients and closed-chest cardiac compression may be ineffective. Rapid delivery of the fetus may improve the response to resuscitative efforts.

11 DESCRIPTION

Ropivacaine hydrochloride injection, USP is a sterile, isotonic solution that contains ropivacaine hydrochloride as the active pharmaceutical ingredient. Ropivacaine hydrochloride is a member of the amino amide class of local anesthetics. Ropivacaine hydrochloride Injection, USP is administered parenterally by infiltration, epidural, and nerve block.

Ropivacaine hydrochloride is chemically described as S-(+)-1-propyl-2',6'-pipecoloylilide hydrochloride. The drug substance is a white crystalline powder, with the following structural formula:



At 25 °C ropivacaine hydrochloride has a solubility of 53.8 mg/mL in water, a distribution ratio between n-octanol and phosphate buffer at pH 7.4 of 14:1 and a pKa of 8.07 in 0.1 M KCl solution. The pKa of ropivacaine is approximately the same as bupivacaine (8.1) and is similar to that of mepivacaine (7.7). However, ropivacaine has an intermediate degree of lipid solubility compared to bupivacaine and mepivacaine.

Ropivacaine hydrochloride injection, USP is a clear, colorless, and preservative-free solution. Each mL contains 2.0 mg, 5.0 mg and 10 mg of Ropivacaine Hydrochloride (equivalent to 2.1 mg, 5.3 mg, 10.6 mg Ropivacaine Hydrochloride monohydrate), and 8.6 mg, 8.0 mg and 7.1 mg of sodium chloride; respectively, and sodium hydroxide and hydrochloric acid as pH adjusters, in water for injection. The pH is adjusted between 4.0 to 6.0. The specific gravity of Ropivacaine hydrochloride injection, USFs solutions range from 1.002 to 1.005 at 25°C.

12. CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Ropivacaine is a member of the amino amide class of local anesthetics and is supplied as the pure S-(+)-enantiomer. Local anesthetics block the generation and the conduction of nerve impulses, presumably by increasing the threshold for electrical excitation in the nerve, by slowing the propagation of the nerve impulses, and by reducing the rate of rise of the action potential. In general, the progression of anesthesia is related to the diameter, myelination and conduction velocity of affected nerve fibers. Clinically, the order of loss of nerve function is as follows: (1) pain, (2) temperature, (3) touch, (4) proprioception, and (5) skeletal muscle tone.

12.2 Pharmacodynamics

Studies in humans have demonstrated that, unlike most other local anesthetics, the presence of epinephrine has no major effect on either the time of onset or the duration of action of ropivacaine. Likewise, addition of epinephrine to Ropivacaine has no effect on limiting systemic absorption of ropivacaine.

Systemic absorption of local anesthetics can produce effects on the central nervous and cardiovascular systems, including: decreased cardiac conduction, excitability, and peripheral vascular resistance have been reported. Toxic blood concentrations depress cardiac conduction and excitability, which may lead to atrioventricular block, ventricular arrhythmias and to cardiac arrest, sometimes resulting in fatalities. In addition, myocardial contractility is depressed and peripheral vasodilation occurs, leading to decreased cardiac output and arterial blood pressure.

Following systemic absorption, local anesthetics can produce central nervous system stimulation, depression or both. Apparent central stimulation is usually manifested as restlessness, tremor and shivering, progressing to convulsions, followed by depression and ultimately to respiratory arrest. However, the local anesthetics have a primary depressant effect on the medulla and on higher centers. The depressed stage may occur without a prior excited stage.

In 2 clinical pharmacology studies (total n=24) ropivacaine and bupivacaine were infused (100 mcg/min) in human volunteers until the appearance of CNS symptoms, e.g., visual or hearing disturbances, perioral numbness, tingling and others. Similar symptoms were seen with both drugs. In 1 study, the mean ± SD maximum tolerated intravenous dose of ropivacaine infused (124 ± 38 mg) was significantly higher than that of bupivacaine (99 ± 30 mg) while in the other study the doses were not different (115 ± 29 mg of bupivacaine). In the latter study, the number of subjects reporting each symptom was similar for both drugs with the exception of muscle twitching which was reported by more subjects with bupivacaine than ropivacaine at comparable intravenous doses. At the end of the infusion, Ropivacaine in both studies caused significantly less depression of cardiac conductivity (less QRS widening) than bupivacaine. Ropivacaine and bupivacaine caused evidence of depression of cardiac conductivity, but there were no changes in cardiac output.

Clinical data in one published article indicate that differences in various pharmacodynamic measures were observed with increasing age. In one study, the upper level of analgesia increased with age, the maximum decrease of mean arterial pressure (MAP) declined with age during the first hour after epidural administration, and the intensity of motor blockade increased with age. However, no pharmacokinetic differences were observed between elderly and younger patients.

In non-clinical pharmacology studies comparing ropivacaine and bupivacaine in several animal species, the cardiac toxicity of ropivacaine was less than that of bupivacaine, although both were considerably more toxic than lidocaine. Arrhythmogenic and cardio-depressant effects were seen in animals at significantly higher doses of ropivacaine than bupivacaine. The incidence of successful resuscitation was not significantly different between the ropivacaine and bupivacaine groups.

12.3 Pharmacokinetics

Absorption

The systemic concentration of ropivacaine is dependent on the total dose and concentration of drug administered, the route of administration, the patient's hemodynamic/circulatory condition, and the vascularity of the administration site.

From the epidural space, ropivacaine shows complete and biphasic absorption. The half-lives of the 2 phases, (mean ± SD) are 14 ± 7 minutes and 4.2 ± 0.9 h, respectively. The slow absorption is the rate limiting factor in the elimination of ropivacaine that explains why the terminal half-life is longer after epidural than after intravenous administration. Ropivacaine shows dose-proportionality up to the highest intravenous dose studied, 80 mg, corresponding to a mean ± SD peak plasma concentration of 1.9 ± 0.3 mcg/mL.

Table 7 Pharmacokinetic (Plasma Concentration-Time) Data from Clinical Trials

Route	Epidural Infusion ^a		Epidural Infusion ^a	Epidural Block [†]	Epidural Block [†]	Plexus Block [‡]	IV Infusion [§]
Dose (mg)	1493 ± 10		1217 ± 277	150	187.5	300	40
n	12	12	11	8	8	10	12
C ₀ (mg/L)	2.4 ± 1*		2.3 ± 1.1*	1.1 ± 0.2	1.6 ± 0.6	2.3 ± 0.8	1.2 ± 0.2*
T _{1/2} (min)	n/a*	n/a		43 ± 14	34 ± 9	54 ± 22	n/a
AUC ₀₋₇₂ (mg·h/L)	135.5 ± 50	145 ± 34	161 ± 90	7.2 ± 2	11.3 ± 4	13 ± 3.3	1.8 ± 0.6
CL (L/h)	11.03	n/a		5.5 ± 2	5 ± 2.6	n/a	21.2 ± 7
T _{1/2} (h)*	5 ± 2.5	5.7 ± 3	6 ± 3	5.7 ± 2	7.1 ± 3	6.8 ± 3.2	1.9 ± 0.5

* Continuous 72 hour epidural infusion after an epidural block with 5 or 10 mg/mL.

† Epidural anesthesia with 7.5 mg/mL (0.75%) for cesarean delivery.

‡ Brachial plexus block with 7.5 mg/mL (0.75%) ropivacaine.

§ 20 minute IV infusion to volunteers (40 mg).

¶ C_{max} measured at the end of infusion (i.e., at 72 hr).

C_{max} measured at the end of infusion (i.e., at 20 minutes).

◆ n/a=not applicable

◆ t_{1/2} is the true terminal elimination half-life. On the other hand, t_{1/2} follows absorption dependent elimination (t_{1/2} (app)) after non-intravenous administration.

In some patients after a 300 mg dose for brachial plexus block, free plasma concentrations of ropivacaine may approach the threshold for CNS toxicity [see Warnings and Precautions (5.7)]. At a dose of greater than 300 mg, for local infiltration, the terminal half-life may be longer (>30 hours).

Distribution

After intravascular infusion, ropivacaine has a steady-state volume of distribution of 41 ± 7 liters. Ropivacaine is 94% protein bound, mainly to α₁-acid glycoprotein. An increase in total plasma concentrations during continuous epidural infusion has been observed, related to a postoperative increase of α₁-acid glycoprotein. Variations in unbound, i.e. pharmacologically active, concentrations have been less than in total plasma concentration. Ropivacaine readily crosses the placenta and equilibrium in regard to unbound concentration will be rapidly reached [see Warnings and Precautions (5) and Use in Specific Population (8.1)].

Metabolism

Ropivacaine is extensively metabolized in the liver, predominantly by aromatic hydroxylation mediated by cytochrome P4501A to 3-hydroxy ropivacaine. After a single IV dose approximately 37% of the total dose is excreted in the urine as both free and conjugated 3-hydroxy ropivacaine. Low concentrations of 3-hydroxy ropivacaine have been found in the plasma. Urinary excretion of the 4 hydroxy ropivacaine, and both the 3-hydroxy N de-alkylated (3-OH-PPX) and 4-hydroxy N de-alkylated (4-OH-PPX) metabolites account for less than 3% of the dose. An additional metabolite, 2-hydroxy-methyl-ropivacaine, has been identified but not quantified in the urine. The N de-alkylated metabolite of ropivacaine PPX and S- OH-ropivacaine are the major metabolites excreted in the urine during epidural infusion. Total PPX concentration in the plasma was about half as that of total ropivacaine; however, mean unbound concentrations of PPX were about 1/3 to 5 times higher than that of unbound ropivacaine following continuous epidural infusion up to 72 hours. Unbound PPX, 3-hydroxy and 4-hydroxy ropivacaine, have a pharmacological activity in animal models less than that of ropivacaine. There is no evidence of in vivo racemization in urine of ropivacaine.

Elimination

The kidney is the main excretory organ for most local anesthetic metabolites. In total, 86% of the ropivacaine dose is excreted in the urine after intravenous administration of which only 1% relates to unchanged drug. After intravenous administration ropivacaine has a mean ± SD total plasma clearance of 387 ± 107 mL/min, an unbound plasma clearance of 7.2 ± 1.6 L/min, and a renal clearance of 1 mL/min. The mean ± SD terminal half-life is 1.8 ± 0.7 h after intravascular administration and 4.2 ± 1 h after epidural administration.

13. NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Long-term studies in animals to evaluate the carcinogenic potential of ropivacaine have not been conducted.

Mutagenesis

Weak mutagenic activity was seen in the mouse lymphoma test. However, ropivacaine was negative in an in vitro Ames assay and an in vivo mouse micronucleus assay.

Impairment of Fertility

No adverse effects on fertility or early embryonic development were reported in a 2-generational reproduction study in which female rats (F₂) were administered subcutaneous doses of 6.3, 12, and 23 mg/kg/day (equivalent to 0.08, 0.15, and 0.29 times the maximum recommended human dose (MRHD) of 770 mg/24 hours for epidural use, respectively, and 0.24, 0.45, and 0.88 times the MRHD of 250 mg for nerve block use, respectively, based on BSA comparisons and a 60 kg human weight) throughout the mating period and pregnancy, partus, and lactation.

13.2 Animal Toxicology and/or Pharmacology

The mean dosages of ropivacaine producing seizures, after intravenous infusion in dogs, nonpregnant and pregnant sheep were 4.9, 6.1 and 5.9 mg/kg (HED: 5.3, 6.6, and 6.4 mg/kg, based on 75 kg sheep weight and 60 kg human weight), respectively. These doses were associated with peak arterial total plasma concentrations of 11.4, 4.3 and 5 mg/mL, respectively.

14. CLINICAL STUDIES

Ropivacaine was studied as a local anesthetic both for surgical anesthesia and for acute pain management [see Dosage and Administration (2)].

The onset, depth and duration of sensory block are, in general, similar to bupivacaine. However, the depth and duration of motor block, in general, are less than that with bupivacaine.

14.1 Epidural Administration in Surgery

There were 25 clinical studies performed in 900 patients to evaluate Ropivacaine hydrochloride epidural injection for general surgery. Ropivacaine hydrochloride was used in doses ranging from 75 to 250 mg. In doses of 100 to 200 mg, the median (1st to 3rd quartile) onset time to achieve a T10 sensory block was 10 (5 to 13) minutes and the median (1st to 3rd quartile) duration at the T10 level was 4 (3 to 5) hours [see Dosage and Administration (2.2)]. Higher doses produced a more profound block with a greater duration of effect.

14.2 Epidural Administration in Cesarean Section

A total of 12 studies were performed with epidural administration of Ropivacaine hydrochloride for cesarean section. Eight of these studies involved 218 patients using the concentration of 5 mg/mL (0.5%) in doses up to 150 mg. Median onset measured at T6 ranged from 11 to 26 minutes. Median duration of sensory block at T6 ranged from 1.7 to 3.2 h, and duration of motor block ranged from 1.4 to 2.8 h. Ropivacaine hydrochloride provided adequate muscle relaxation for surgery in all cases.

In addition, 4 active controlled studies for cesarean section were performed in 264 patients at a concentration of 7.5 mg/mL (0.75%) in doses up to 187.5 mg. Median onset measured at T6 ranged from 4 to 15 minutes. Seventy-seven to 96% of Ropivacaine hydrochloride-exposed patients reported no pain at delivery. Some patients received other anesthetic, analgesic, or sedative modalities during the course of the operative procedure.

14.3 Epidural Administration in Labor and Delivery

A total of 9 double-blind clinical studies, involving 240 patients were performed to evaluate Ropivacaine hydrochloride for epidural block for management of labor pain. When administered in doses up to 278 mg as intermittent injections or as a continuous infusion, Ropivacaine hydrochloride produced adequate pain relief. A prospective meta-analysis on 6 of these studies provided detailed evaluation of the delivered newborns and showed no difference in clinical outcomes compared to bupivacaine. There were significantly fewer instrumental deliveries in mothers receiving ropivacaine as compared to bupivacaine.

Table 8 Labor and Delivery Meta-analysis: Mode of Delivery

Delivery Mode	Ropivacaine hydrochloride n = 199		Bupivacaine n = 188	
	n	%	n	%
Spontaneous Vaginal	116	58	92	49
Vacuum Extractor	26		33	
		32*		34§
Forceps	28		42	
Cesarean Section	29	15	21	11

*p=0.004 versus bupivacaine

14.4 Epidural Administration in Postoperative Pain Management

There were 8 clinical studies performed in 382 patients to evaluate Ropivacaine hydrochloride 2 mg/mL (0.2%) for postoperative pain management after upper and lower abdominal surgery and after orthopedic surgery. The studies utilized intravascular morphine via PCA as a rescue medication and quantified as an efficacy variable.

Epidural anesthesia with Ropivacaine hydrochloride 5 mg/mL (0.5%) was used intraoperatively for each of these procedures prior to initiation of postoperative Ropivacaine hydrochloride. The incidence and intensity of the motor block were dependent on the dose-rate of Ropivacaine hydrochloride and the site of injection. Cumulative doses of up to 770 mg of ropivacaine were administered over 24 hours (intraoperative block plus postoperative continuous infusion). The overall quality of pain relief, as judged by the patients, in the ropivacaine groups was rated as good or excellent (73% to 100%). The frequency of motor block was greatest at 4 hours and decreased during the infusion period in all groups. At least 80% of patients in the upper and lower abdominal studies and 42% in the orthopedic studies had no motor block at the end of the 21-hour infusion period. Sensory block was also dose rate dependent and a decrease in spread was observed during the infusion period.

A double-blind, randomized, clinical trial compared lumbar epidural infusion of Ropivacaine hydrochloride (n=26) and bupivacaine (n=26) at 2 mg/mL (8 mL/h), for 24 hours after knee replacement. In this study, the pain scores were higher in the Ropivacaine hydrochloride group, but the incidence and the intensity of motor block were lower.

Continuous epidural infusion of Ropivacaine hydrochloride 2 mg/mL (0.2%) during up to 72 hours for postoperative pain management after major abdominal surgery was studied in 2 multicenter, double-blind studies. A total of 391 patients received a low thoracic epidural catheter, and Ropivacaine hydrochloride 7.5 mg/L (0.75%) was given for surgery in combination with GA. Postoperatively, Ropivacaine (0.2%), 4 to 14 mL/h, alone or with fentanyl 1, 2, or 4 mcg/mL, was infused through the epidural catheter and adjusted according to the patient's needs. These studies support the use of Ropivacaine hydrochloride 2 mg/mL (0.2%) for epidural infusion at 6 to 14 mL/h (12 to 28 mg) for up to 72 hours and demonstrated adequate analgesia with only slight and nonprogressive motor block in cases of moderate to severe postoperative pain.

Clinical studies with 2 mg/mL (0.2%) Ropivacaine hydrochloride have demonstrated that infusion rates of 6 to 14 mL (12 to 28 mg) per hour provide adequate analgesia with nonprogressive motor block in cases of moderate to severe postoperative pain. In these studies, this technique resulted in a significant reduction in patients' morphine rescue dose requirement. Clinical experience supports the use of Ropivacaine hydrochloride injection epidural infusions for up to 72 hours.

14.5 Peripheral Nerve Block

Ropivacaine hydrochloride, 5 mg/mL (0.5%), was evaluated for its ability to provide anesthesia for surgery using the techniques of Peripheral Nerve Block. There were 13 studies performed including a series of 4 pharmacodynamic and pharmacokinetic studies performed on minor nerve blocks. From these, 235 Ropivacaine hydrochloride-treated patients were available for efficacy. Ropivacaine hydrochloride was used in doses up to 275 mg. When used for brachial plexus block, onset depended on technique used. Suprascapular blocks were consistently more successful than axillary blocks. The median onset of sensory block (anesthesia) produced by ropivacaine 0.5% via axillary block ranged from 10 minutes (medial brachial cutaneous nerve) to 45 minutes (musculocutaneous nerve). Median duration ranged from 3.7 hours (medial brachial cutaneous nerve) to 8.7 hours (ulnar nerve). The 5 mg/mL (0.5%) Ropivacaine hydrochloride solution gave success rates from 56% to 98% for axillary blocks, compared with 52% for suprascapular blocks.

In addition, Ropivacaine hydrochloride, 7.5 mg/mL (0.75%), was evaluated in 98 Ropivacaine hydrochloride-treated patients, in 2 double-blind studies, performed to provide anesthesia for surgery using the techniques of Brachial Plexus Block. Ropivacaine hydrochloride 7.5 mg/mL was compared to bupivacaine 5 mg/mL. In 1 study, patients underwent axillary brachial plexus block using injections of 40 mL (300 mg) of Ropivacaine hydrochloride, 7.5 mg/mL (0.75%) or 40 mL injections of bupivacaine, 5 mg/mL (200 mg). In a second study, patients underwent subclavian perivascular brachial plexus block using 30 mL (225 mg) of Ropivacaine hydrochloride injection, 7.5 mg/mL (0.75%) or 30 mL of bupivacaine 5 mg/mL (150 mg). There was no significant difference between the Ropivacaine hydrochloride and bupivacaine groups in either study with regard to onset of anesthesia, duration of sensory blockade, or duration of anesthesia.

The median duration of anesthesia varied between 11.4 and 14.4 hours with both techniques. In one study, using the axillary technique, the quality of analgesia and muscle relaxation in the Ropivacaine hydrochloride group was judged to be significantly superior to bupivacaine by both investigator and surgeon. However, using the subclavian perivascular technique, no statistically significant difference was found in the quality of analgesia and muscle relaxation as judged by both the investigator and surgeon. The use of Ropivacaine hydrochloride injection 7.5 mg/mL for block of the brachial plexus via either the subclavian perivascular approach using 30 mL (225 mg) or via the axillary approach using 40 mL (300 mg) both provided effective and reliable anesthesia.

14.6 Local Infiltration

A total of 7 clinical studies were performed to evaluate the local infiltration of Ropivacaine hydrochloride injection to produce anesthesia for surgery and analgesia in postoperative pain management. In these studies 297 patients who received Ropivacaine hydrochloride in doses up to 200 mg (concentrations up to 5 mg/mL, 0.5%) were available for efficacy. With infiltration of 100 to 200 mg Ropivacaine hydrochloride, the time to first request for analgesic was 2 to 6 hours. When compared to placebo, Ropivacaine hydrochloride produced lower pain scores and a reduction of analgesic consumption.

16. How Supplied/Storage and Handling

Ropivacaine hydrochloride injection USP is a clear, colorless, and preservative-free solution, available in single-dose containers in 2 mg/mL (0.2%), 5 mg/mL (0.5%) and 10 mg/mL (1%) concentrations.

Storage

Solutions should be stored at 20 °C to 25 °C (68 °F to 77 °F) [See USP Controlled Room Temperature].

Ropivacaine Hydrochloride Injection, USP Single-Dose Vials

NDC No.	Strength	Vial Size
65145-107-10	0.2% - 40 mg/20 mL (2 mg/mL)	20 mL single dose vial, in packages of 10.
65145-108-10	0.5% - 100 mg/20 mL (5 mg/mL)	20 mL single dose vial, in packages of 10.
65145-109-10	0.5% - 150 mg/30 mL (5 mg/mL)	30 mL single dose vial, in packages of 10.
65145-110-10	1% - 200 mg/20 mL (10mg/mL)	20 mL single dose vial, in packages of 10.

Ropivacaine hydrochloride injection single-dose, ready-to-use, polypropylene flexible bags overwrapped with aluminum pouch. The flexible bag container is not made with natural rubber latex or polyvinyl chloride (PVC), Non-DEHP.

NDC No	Strength	Size
NDC 65145-165-01	200 mg/100 mL (2 mg/mL)	100 mL infusion bag
NDC 65145-166-01	400 mg/200 mL (2 mg/mL)	250 mL infusion bag

Ropivacaine hydrochloride container closure is not made with natural rubber latex.

These products are intended for single-dose and are free from preservatives. Discard Unused Portion.

17. PATIENT COUNSELING INFORMATION

17.1 Information for Patients and Caregivers

When appropriate, patients should be informed in advance that they may experience temporary loss of sensation and motor activity in the anesthetized part of the body following proper administration of lumbar epidural anesthesia. Also, when appropriate, the physician should discuss other information including adverse reactions in the Ropivacaine hydrochloride package insert.

Inform patients that use of local anesthetics may cause methemoglobinemia, a serious condition that must be treated promptly. Advise patients or caregivers to seek immediate medical attention if they or someone in their care experience the following signs or symptoms: pale, gray, or blue colored skin (cyanosis); headache; rapid heart rate; shortness of breath; lightheadedness; or fatigue.



Made in India

Distributed by :

Caplin Steriles USA Inc,

Hamilton, NJ 080619,

Code: TNDrogsTM00006405